Complementary Feed For Poultry



MIXTRAL

' Hydration management

Approved to use UAB for organic farming.



- Promote drinking behaviour and consumption of drinking water (starting period, transfer, hydration)
- Contribute to manage urinary problems
- Support kidney functions





Main components


- Plants: Blackcurrant, Cranberry, Orthosiphon, Meadowsweet
- Magnesium chloride
- ▶ Flavouring substances



1 to 3 ml per liter of water for 1 to 3 days.

▶ Starting period & transfer (layers)

Preceded by BACI 50 following the estimation of a possible colibacillary disorder



After a renal elimination antibiotic

After a complicated sanitary passage that required an intervention with an antibiotic molecule



^{*} For more details on the products mentioned, please refer to their commercial sheet

Résultats

Marocco trial 2017 - Pullets for lay during starting period

	Chick weight		
	MIXTRAL IMUNEO Protocol	Control	
D 0	33 g	33 g	
D 7	69 g	64 g	+5g=+8%
D 15	124 g	117 g	+7g=+6%

8	000
	8

MIXTRAL D 0 et D +1 / IMUNEO D +2 à D +5

Mortality			
MIXTRAL IMUNEO Protocol	Control		
-	-		
0,34 %	0,37 %		
0,22 %	0,31 %		
0,56 %	0,68 %		

- 12 pts = - 17 %

